


Sexually Transmitted Disease Profile

Grant County 2004



Summary

This report describes the sexually transmitted disease burden in Grant County. Primary emphasis is placed on chlamydia and gonorrhea since they are the most frequently reported STDs in Washington State. The 2004 incidence rates by age and sex for gonorrhea and chlamydia are presented. The report concludes with a presentation of which providers in your county reported STDs.

Contents

County STD Trends

Table 1: Washington State Reportable Sexually Transmitted Diseases, 2004	1
--	---

Chlamydia

Figure 1: Chlamydia Incidence Rates, by Age and Gender, 2004.....	2
Figure 2: Chlamydia Cases by Age (13-19) and Gender, 2004.....	3
Table 2: Chlamydia Repeater Infections, 2004	3
Table 3: Chlamydia Asymptomatic Infections, 2004	4

Gonorrhea

Figure 3: Gonorrhea Incidence Rates, by Age and Gender, 2004	5
Table 4: Gonorrhea Symptomatic Infections, 2004.....	5

Conclusion

Table 5: Reported Cases of Gonorrhea and Chlamydia by Provider Type, 2004	6
Health People 2010 National Objective for Chlamydia	6
Health People 2010 National Objective for Gonorrhea.....	7

Appendix

A. Data Sources, Analyses, and Limitations	8
--	---

Grant County STD Disease Trends

Table 1: **Washington State** Reportable Sexually Transmitted Diseases, Grant County, 2004

Disease	2003 Grant County Cases	2004 Grant County Cases	2004 Grant County Rate ^λ (per 100,000)	2004 Washington State Rate ^λ (per 100,000)
Chlamydia	216	234	299	286
Gonorrhea	13	15	19	46
Early Syphilis	0	0	-	3.3
Congenital Syphilis	0	0	-	0.0 (live births)
Late/Late Latent Syphilis	1	0	*	2.2
Herpes (initial infection)	15	30	38	35
GI/LGV/Chancroid**	0	0	-	0.0
HIV cases**	1	2		
AIDS cases**	1	2		
TOTAL (excluding HIV/AIDS cases)	245	279	356	372

^λ Denominator estimates for the calculation of incidence rates from Washington State Adjusted Population Estimates, OFM, February 2004.

* Rates cannot be calculated for years with fewer than five cases.

** See Appendix A for explanation of disease acronyms.

In 2004, Grant County experienced an increase from 2003 in its combined reportable STD cases. With 279 cases of STDs (excluding HIV/AIDS cases¹) in 2004, the incidence rate for all STDs was 356 per 100,000 persons. This is 4% less than the 372 per 100,000 combined reportable STD rate for Washington State in 2004. Grant County reported no cases of congenital syphilis or GI/LGV/Chancroid in 2004.


2004 compared to 2003:

- Chlamydia had an 8% increase in reported cases (234 vs. 216).
- Gonorrhea had a 15% increase in reported cases (15 vs. 13).
- Late/Late Latent Syphilis had a 100% decrease in reported cases (0 vs. 1).
- Initial infection herpes had a 100% increase in reported cases (30 vs. 15).

¹ Complete information on the HIV/AIDS epidemic in Washington can be found in [Washington State HIV/AIDS Surveillance Report](#), Washington State Department of Health, IDRH Assessment Unit.

Chlamydia

Figure 1: **Chlamydia** Incidence Rates by Age and Gender, Grant County, 2004



Female Rate	0	*	2,404	2,395	1,113	475	*	*
Male Rate	0	0	360	872	313	*	*	*
Female Cases	0	2	76	61	26	11	4	1
Male Cases	0	0	13	26	8	3	1	1

[^] Denominator estimates for the calculation of incidence rates from Washington State Adjusted Population Estimates, OFM, February 2004. Incidence rates rounded to the nearest whole number.


* Rates cannot be calculated for ages with fewer than five cases.

In 2004 the female chlamydia incidence rate was highest among the 15-19 year old age group in Grant County, at 2,404 cases per 100,000. Chlamydia incidence among females progressively declined with increasing age. Among the men in Grant County, the 2004 chlamydia incidence rate was highest among 20-24 year olds at 872 per 100,000 then declined with increasing age.

Women are preferentially screened for chlamydia. Because active case-finding is preferentially limited to women, the incidence of chlamydia in men may be under-reported by comparison. Caution should be used in interpreting comparisons of chlamydia rates between genders.

The 2002 STD Treatment Guidelines from CDC recommend that all women diagnosed with chlamydia be re-screened three to four months after treatment. This was suggested because of the high prevalence of chlamydia found in women diagnosed with the disease in the preceding months, presumably as a result of re-infection.

Figure 2: **Chlamydia** Cases by Age (13 – 19) and Gender, Grant County, 2004



Repeater Infections (Persons having more than one infection in a 12-month period.)

Recurrent infection is common and associated with increased risk of pelvic inflammatory disease (PID) and other serious outcomes. Data suggest that young age and incomplete therapy increase the risk for persistent/recurrent infection. Studies also suggest that women's current male sex partners are not receiving treatment for chlamydia and that women are being re-infected by resuming sex with previous (and infected) sex partners. Careful interviewing and prompt, concurrent treatment of all partners is important. People should be coached to ask health care providers for re-screening.

Table 2: **Chlamydia** Repeater Infections, Grant County, 2004.

	MALE	FEMALE	TOTAL
Reported Cases	52	182	234
Repeaters Identified	1	12	13
% Repeaters	1.9%	6.6%	5.6%

Asymptomatic Infection

STD infections often lack any signs and symptoms. Routine screening and treatment is essential to prevent serious complications that may not appear until long after infection. Screening all sexually active adolescents (19 years and younger) during sports physicals and routine office visits should be done even if symptoms are not present. Screening women and men aged 20-25 is also suggested, particularly those who have new or multiple sex partners. Women who are

pregnant, have sex partners infected with chlamydia, have mucopurulent cervicitis or are planning an IUD insertion should also be screened. Careful interviewing and treatment of all partners is important.

Table 3: Reported Cases of **Chlamydia** by Diagnostic Category, Grant County, 2004.

Diagnosis	Private		Public		Total		Total Cases
	Male	Female	Male	Female	Male	Female	
Asymptomatic	14	113	9	11	23	124	147
Symptomatic-Uncomplicated	26	54	3	0	29	54	83
Pelvic Inflammatory Disease	0	3	0	0	0	3	3
Other	0	0	0	0	0	0	0
Unknown	0	1	0	0	0	1	1
TOTAL	40	171	12	11	52	182	234

Gonorrhea

Grant County gonorrhea rates for 2004 by age groups cannot be calculated because all age groups have less than five cases.

Figure 3: **Gonorrhea** Incidence Rates, by Age and Gender, Grant County 2004

	0-9	10-14	15-19	20-24	25-29	30-34	35-39	40+
Female Rate	0	0	*	*	*	0	0	*
Male Rate	0	0	0	*	*	0	0	0
Female Cases	0	0	3	2	3	0	0	1
Male Cases	0	0	0	4	2	0	0	0

[^] Denominator estimates for the calculation of incidence rates from Washington State Adjusted Population Estimates, OFM, February 2004. Incidence rates rounded to the nearest whole number.

* Rates cannot be calculated for years with fewer than five cases.

In Washington State the reported rate of gonorrhea incidence in 2004 was 46/100,000, a slight increase from the 2003 rate. Statewide, the greatest incidence of disease among females was among 15-19 year olds (198/100,000), while for males the burden of disease is distributed more evenly among those older. Males had a higher gonorrhea rate (51/100,000) than females (40/100,000). A major factor contributing to the distribution of gonorrhea incidence in different age groups among men or women is a documented outbreak among men who have sex with men (MSM) whose median reported age was 30.

Findings from the Gonococcal Isolate Surveillance Project (GISP) in Seattle have indicated that Washington State is now an area with increased prevalence of quinolone-resistant *Neisseria gonorrhoeae* (QRNG). Based on these findings, the Washington State Department of Health recommends that health care providers in the state should no longer use fluoroquinolones (ciprofloxacin, levofloxacin and ofloxacin) as first line therapy for gonorrhea. The antibiotics of choice are ceftriaxone (Rocephin™) or cefpodoxime (Vantin™) followed with either azithromycin or doxycycline to treat possible coexisting chlamydial infection.

Because most gonorrhea cases are symptomatic and seek medical care, reported cases are considered to be an accurate reflection of true disease incidence in the overall population. Providers in Washington State who reported gonorrhea cases in 2004 indicated that 83% of the men were symptomatic for gonorrhea; 47% of the women were symptomatic.

Table 4: Reported Cases of **Gonorrhea** by Diagnostic Category, Grant County, 2004.

Diagnosis	Private		Public		Total		Total Cases
	Male	Female	Male	Female	Male	Female	
Asymptomatic	1	6	0	0	1	6	7
Symptomatic-Uncomplicated	5	3	0	0	5	3	8
Pelvic Inflammatory Disease	0	0	0	0	0	0	0
Other	0	0	0	0	0	0	0
Unknown	0	0	0	0	0	0	0
TOTAL	6	9	0	0	6	9	15

Conclusion

Table 5: Reported Cases of Chlamydia and Gonorrhea by Provider Type, Grant County, 2004

Provider Type	Chlamydia			Gonorrhea		
	No. of Providers	No. of Cases	Percent of Total Cases	No. of Providers	No. of Cases	Percent of Total Cases
Alcohol/Substance Abuse						
Blood Bank/Plasma Center						
Community Health Center						
Emergency Care (excl. hosp.)						
Family Planning	6	48	21%	1	2	13%
Health Plan/HMOs						
HIV/AIDS						
Hospitals	1	2	1%			
Indian Health	2	3	1%			
Jail/Correction/Detention	1	1	0%			
Job Corps	1	19	8%			
Migrant Health	2	45	19%	2	6	40%
Military						
Neighborhood Health						
OB/GYN	5	8	3%			
Other	20	105	45%	3	7	47%
Private Physicians	1	3	1%			
Reproductive Health						
STD Clinics						
Student Health						
TOTAL	39	234	100%	6	15	100%

In Grant County, the other providers reported the highest number of chlamydia cases. These providers reported 45% of the total. Family planning reported the second highest number of chlamydia cases (21%). Gonorrhea cases were most frequently reported by other providers (47%).

In 2004, 89% of chlamydia and 100% of gonorrhea cases indicating treatment date by health care providers in Grant County were reported within 30 days to the STD Washington State Morbidity System. Treatment date for cases reported by providers in Grant County, were missing on 3% of chlamydia and 7% of gonorrhea cases in 2004.

The **Healthy People 2010** national objectives for chlamydia incidence are:

Females aged 15-24 attending family planning clinics: 3%.

There is one (1) Region X Infertility Prevention Project (IPP) family planning clinic in Grant County. The 2004 positivity rate for females was:

Sites	<u>Male</u>			<u>Female</u>		
	No. of Tests	No. of Pos.	Percent Pos.	No. of Tests	No. of Pos.	Percent Pos.
FP Services of Grant Co.	56	12	21.4	517	27	5.2

Females aged 15-24 attending STD clinics: 3%.

Males aged 15-24 attending STD clinics: 3%.

There are zero (0) Region X IPP STD/reproductive health clinics in Grant County.

Other Region X Infertility Prevention Project Sites in Grant County include:

Sites	<u>Male</u>			<u>Female</u>		
	No. of Tests	No. of Pos.	Percent Pos.	No. of Tests	No. of Pos.	Percent Pos.
Clinic at Royal City	3	0	0.0	21	1	4.8
Mattawa Medical Clinic	10	1	10.0	23	2	8.7

See attachment A for Region X IPP screening criteria.

The **2010 Healthy People** national objective for gonorrhea incidence is 19 cases per 100,000.

Grant County has met this goal with the 2004 rate of 19 cases per 100,000.

The Aptima test used to diagnose chlamydia is a combined test that will also diagnose gonorrhea. Gonorrhea positives from the Region X IPP sites include:

Family Planning Site	<u>Male</u>			<u>Female</u>		
	No. of Tests	No. of Pos.	Percent Pos.	No. of Tests	No. of Pos.	Percent Pos.
FP Services of Grant Co.	56	1	1.8	513	1	0.2

Other Site	<u>Male</u>			<u>Female</u>		
	No. of Tests	No. of Pos.	Percent Pos.	No. of Tests	No. of Pos.	Percent Pos.
Clinic at Royal City	3	0	0.0	10	0	0.0
Mattawa Medical Clinic	10	0	0.0	26	1	3.8

Appendix A: Data Sources, Analyses and Limitations

Cases: The number of cases identified and submitted by providers to local health jurisdictions and forwarded to the Washington State Department of Health, Office of Infectious Disease and Reproductive Health, STD/TB Services.

Population: Denominator population estimates for incidence rates are from Washington State Adjusted Population Estimates, Office of Financial Management (OFM), February 2004.

Incidence Rates: Incidence rates are calculated as the number of new episodes of a disease (not persons) in a given year divided by the total population (age and sex appropriate) for that year, expressed as a rate per 100,000. Incidence rates allow comparisons between two or more populations by standardizing the denominator and are the most appropriate statistic to use when investigating differences between groups. Rates should not be calculated for incident case totals fewer than five because the rates are unstable.

Data Reporting: Gonorrhea, chlamydia, syphilis, and herpes (initial infection) are reportable diseases to the local health jurisdictions and forwarded to the Department of Health. To be reported and included in surveillance data, disease definition must be met.

Disease Definitions:

- AIDS - Acquired Immunodeficiency Syndrome is the advanced stage of HIV-disease in humans and is characterized by severe suppression of immune response. Persons with AIDS are at risk for increased susceptibility to opportunistic infections, degradation of major organ systems and eventual death.
- Chancroid - a STD characterized by painful genital ulceration and inflammatory inguinal adenopathy caused by the bacterium *Haemophilus ducreyi*.
- Chlamydia- isolation of *Chlamydia trachomatis* from a clinical specimen by culture or non-culture methods that detect chlamydia antigen or genetic material.
- Gonorrhea - isolation of *Neisseria gonorrhoeae* from a clinical specimen by culture or non-culture methods or observation of Gram-negative intracellular diplococci in urethral or endocervical smears.
- Granuloma Inguinale (GI) - a slowly progressive ulcerative disease of the skin and lymphatics of the genital and perianal area.
- Herpes Simplex (initial infection only) - diagnostic criteria for reporting can be made through clinical observation of typical lesions and/or laboratory confirmation.
- HIV - Human Immunodeficiency Virus is a retrovirus causing HIV disease and AIDS in humans. This pathogen is transmitted from person to person through unprotected sexual contact, sharing of injection equipment and transfusion/transplantation with infected blood or tissue.
- Lymphogranuloma Venereum (LGV) - characterized by genital lesions, suppurative regional lymphadenopathy, or hemorrhagic proctitis, caused by the L1, L2 and L3 serovars of *Chlamydia trachomatis*.

- Syphilis - a complex sexual transmitted disease with a highly variable clinical course. See CDC guidelines for surveillance definition.

The diagnosing practitioner is responsible for providing the case information which includes patient demographics, source of diagnosis, limited clinical information including site of infection and treatment, and date of diagnosis.

Data Strengths: Sexually transmitted disease data may provide more timely information on behavioral trends in the community than diseases with similar modes of transmission particularly HIV/AIDS. There is a high level of participation in the STD surveillance system by private providers of STD services.

Data Limitations: Clinically diagnosed cases of STDs (without laboratory confirmation) may be missed through this surveillance system. Depending upon diagnosing practices, completeness of reporting may vary by source of health care.

Data Biases: Biases could exist in the data due to under-reporting, inability of certain populations to access medical services, error in laboratory reporting, or differential reporting or screening by disease and source of care. However, it is assumed that the number of cases that would fall into these categories is small and normally distributed, thus not significantly impacting the calculated STD rates.

Assumptions: It is assumed that the cases reported from year to year are independent of each other. One violation of this assumption could be if a person who has an STD one year is more likely to have an STD the following year. Also, repeat episodes of the same STD by the same person are not excluded from the numerator count; it is felt that these numbers are not large enough to significantly impact the calculated incidence rates. Finally, we have assumed that all rates follow a chi-square distribution.

Region X IPP screening criteria that are used at the 140 IPP clinics are as follows:

- Sexually active women 24 years and younger;
- Pregnant women;
- Women with mucopurulent cervicitis, cervical friability, or ectopy with inflammation or edema;
- Women with pelvic inflammatory disease (PID);
- Women planning to receive an intrauterine device;
- Women with a symptomatic sex partner;
- Women diagnosed with CT in the last 12 months; and
- Sex partners of persons with chlamydial infection.